



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 15, 2026 – 10:15 AM JST

PDB ID : 24FN / pdb_000024fn
Title : Dextran glucosidase SmDG E236S in complex with isomaltotriose
Authors : Saburi, W.; Ose, T.; Yao, M.; Mori, H.
Deposited on : 2026-03-02
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	543	 88% 11% .
1	B	543	 87% 12% .
1	C	543	 86% 12% .
1	D	543	 87% 12% .
2	E	2	 100%
2	F	2	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	2	 100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 18017 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucan 1,6-alpha-glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	536	Total	C	N	O	S	0	0	0
			4370	2794	729	825	22			
1	B	536	Total	C	N	O	S	0	0	0
			4369	2794	728	825	22			
1	C	536	Total	C	N	O	S	0	0	0
			4370	2794	729	825	22			
1	D	536	Total	C	N	O	S	0	0	0
			4370	2794	729	825	22			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	72	ALA	THR	conflict	UNP Q99040
A	236	SER	GLU	engineered mutation	UNP Q99040
A	536	LEU	-	expression tag	UNP Q99040
A	537	GLU	-	expression tag	UNP Q99040
A	538	HIS	-	expression tag	UNP Q99040
A	539	HIS	-	expression tag	UNP Q99040
A	540	HIS	-	expression tag	UNP Q99040
A	541	HIS	-	expression tag	UNP Q99040
A	542	HIS	-	expression tag	UNP Q99040
A	543	HIS	-	expression tag	UNP Q99040
B	72	ALA	THR	conflict	UNP Q99040
B	236	SER	GLU	engineered mutation	UNP Q99040
B	536	LEU	-	expression tag	UNP Q99040
B	537	GLU	-	expression tag	UNP Q99040
B	538	HIS	-	expression tag	UNP Q99040
B	539	HIS	-	expression tag	UNP Q99040
B	540	HIS	-	expression tag	UNP Q99040
B	541	HIS	-	expression tag	UNP Q99040
B	542	HIS	-	expression tag	UNP Q99040
B	543	HIS	-	expression tag	UNP Q99040
C	72	ALA	THR	conflict	UNP Q99040

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	236	SER	GLU	engineered mutation	UNP Q99040
C	536	LEU	-	expression tag	UNP Q99040
C	537	GLU	-	expression tag	UNP Q99040
C	538	HIS	-	expression tag	UNP Q99040
C	539	HIS	-	expression tag	UNP Q99040
C	540	HIS	-	expression tag	UNP Q99040
C	541	HIS	-	expression tag	UNP Q99040
C	542	HIS	-	expression tag	UNP Q99040
C	543	HIS	-	expression tag	UNP Q99040
D	72	ALA	THR	conflict	UNP Q99040
D	236	SER	GLU	engineered mutation	UNP Q99040
D	536	LEU	-	expression tag	UNP Q99040
D	537	GLU	-	expression tag	UNP Q99040
D	538	HIS	-	expression tag	UNP Q99040
D	539	HIS	-	expression tag	UNP Q99040
D	540	HIS	-	expression tag	UNP Q99040
D	541	HIS	-	expression tag	UNP Q99040
D	542	HIS	-	expression tag	UNP Q99040
D	543	HIS	-	expression tag	UNP Q99040

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	2	Total	C	O	0	0	0
			22	12	10			
2	F	2	Total	C	O	0	0	0
			22	12	10			
2	G	2	Total	C	O	0	0	0
			22	12	10			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		
3	B	2	Total	Ca	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	2	Total	Ca	0	0
			2	2		
3	D	2	Total	Ca	0	0
			2	2		

- Molecule 4 is alpha-D-glucopyranose (CCD ID: GLC) (formula: C₆H₁₂O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			11	6	5		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	111	Total	O	0	0
			111	111		
5	B	109	Total	O	0	0
			109	109		
5	C	113	Total	O	0	0
			113	113		
5	D	120	Total	O	0	0
			120	120		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucan 1,6-alpha-glucosidase




- Molecule 1: Glucan 1,6-alpha-glucosidase



- Molecule 1: Glucan 1,6-alpha-glucosidase




- Molecule 1: Glucan 1,6- α -glucosidase

Chain D:  87% 12%



- Molecule 2: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose

Chain E:  100%



- Molecule 2: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose

Chain F: 100%



- Molecule 2: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose

Chain G: 100%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	134.47Å 73.08Å 134.80Å 90.00° 102.53° 90.00°	Depositor
Resolution (Å)	48.90 – 2.40 48.90 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.90-2.40) 99.7 (48.90-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.39Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.186 , 0.253 0.188 , 0.254	Depositor DCC
R_{free} test set	2012 reflections (2.01%)	wwPDB-VP
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.613	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.098 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18017	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.33 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.5717e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/4486	0.52	0/6087
1	B	0.34	0/4485	0.52	0/6085
1	C	0.35	0/4486	0.51	0/6087
1	D	0.35	0/4486	0.52	0/6087
All	All	0.34	0/17943	0.52	0/24346

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4370	0	4208	34	0
1	B	4369	0	4208	40	0
1	C	4370	0	4210	44	0
1	D	4370	0	4210	40	0
2	E	22	0	18	0	0
2	F	22	0	19	0	0
2	G	22	0	19	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	D	11	0	10	0	0
5	A	111	0	0	1	0
5	B	109	0	0	1	0
5	C	113	0	0	2	0
5	D	120	0	0	1	0
All	All	18017	0	16902	157	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 157 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:484:GLU:O	1:D:485:GLU:HG3	1.70	0.90
1:D:366:GLU:HG2	1:D:424:PRO:HG3	1.70	0.74
1:B:337:ILE:HD13	1:B:451:LEU:HD11	1.73	0.71
1:B:280:LYS:H	1:B:280:LYS:HD2	1.53	0.70
1:C:329:GLU:HG3	1:C:444:ILE:HG13	1.74	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	534/543 (98%)	518 (97%)	16 (3%)	0	100	100
1	B	534/543 (98%)	513 (96%)	21 (4%)	0	100	100
1	C	534/543 (98%)	514 (96%)	19 (4%)	1 (0%)	43	58
1	D	534/543 (98%)	511 (96%)	23 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	2136/2172 (98%)	2056 (96%)	79 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	195	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	470/477 (98%)	469 (100%)	1 (0%)	87	94
1	B	470/477 (98%)	469 (100%)	1 (0%)	87	94
1	C	470/477 (98%)	469 (100%)	1 (0%)	87	94
1	D	470/477 (98%)	469 (100%)	1 (0%)	87	94
All	All	1880/1908 (98%)	1876 (100%)	4 (0%)	87	94

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	310	ASN
1	B	310	ASN
1	C	310	ASN
1	D	310	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	526	GLN
1	D	292	ASN
1	D	365	ASN
1	D	174	GLN
1	B	458	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	E	1	2	11,11,12	0.86	0	15,15,17	2.51	5 (33%)
2	GLC	E	2	2	11,11,12	0.63	0	15,15,17	3.04	6 (40%)
2	GLC	F	1	2	11,11,12	0.70	0	15,15,17	1.10	2 (13%)
2	GLC	F	2	2	11,11,12	0.68	0	15,15,17	2.07	5 (33%)
2	GLC	G	1	2	11,11,12	0.57	0	15,15,17	2.13	4 (26%)
2	GLC	G	2	2	11,11,12	0.56	0	15,15,17	2.05	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	1	2	-	2/2/19/22	0/1/1/1
2	GLC	E	2	2	-	0/2/19/22	0/1/1/1
2	GLC	F	1	2	-	2/2/19/22	0/1/1/1
2	GLC	F	2	2	-	0/2/19/22	0/1/1/1
2	GLC	G	1	2	-	2/2/19/22	0/1/1/1
2	GLC	G	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2	GLC	O5-C1-C2	-8.36	97.87	110.77
2	E	1	GLC	C1-C2-C3	7.08	118.37	109.67
2	F	2	GLC	O5-C1-C2	-5.50	102.28	110.77
2	G	2	GLC	O5-C1-C2	-5.16	102.80	110.77
2	G	1	GLC	C1-O5-C5	4.29	118.00	112.19

There are no chirality outliers.



5 of 6 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	1	GLC	O5-C5-C6-O6
2	G	1	GLC	C4-C5-C6-O6
2	G	1	GLC	O5-C5-C6-O6
2	E	1	GLC	C4-C5-C6-O6
2	F	1	GLC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 8 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GLC	D	601	-	11,11,12	0.76	0	15,15,17	2.46	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GLC	D	601	-	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	601	GLC	O2-C2-C3	4.88	119.92	110.14
4	D	601	GLC	O5-C1-C2	-4.87	103.26	110.77
4	D	601	GLC	C1-C2-C3	-4.48	104.16	109.67
4	D	601	GLC	C3-C4-C5	2.18	114.12	110.24
4	D	601	GLC	C1-O5-C5	2.11	115.05	112.19


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	536/543 (98%)	-0.20	1 (0%) 91 89	21, 31, 47, 64	0
1	B	536/543 (98%)	-0.30	1 (0%) 91 89	20, 30, 45, 58	0
1	C	536/543 (98%)	-0.31	1 (0%) 91 89	20, 30, 46, 65	0
1	D	536/543 (98%)	-0.32	2 (0%) 88 86	20, 30, 45, 66	0
All	All	2144/2172 (98%)	-0.28	5 (0%) 91 89	20, 30, 46, 66	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	TRP	3.6
1	D	238	TRP	2.5
1	C	238	TRP	2.3
1	B	365	ASN	2.1
1	D	465	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	E	1	11/12	0.79	0.14	31,43,54,65	0
2	GLC	F	1	11/12	0.90	0.10	33,36,41,49	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLC	G	1	11/12	0.90	0.09	34,36,41,54	0
2	GLC	E	2	11/12	0.94	0.08	23,31,34,35	0
2	GLC	F	2	11/12	0.95	0.07	28,31,34,37	0
2	GLC	G	2	11/12	0.95	0.07	25,27,35,36	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GLC	D	601	11/12	0.90	0.13	24,30,36,37	0
3	CA	D	603	1/1	0.91	0.07	58,58,58,58	0
3	CA	A	602	1/1	0.91	0.08	51,51,51,51	0
3	CA	C	602	1/1	0.93	0.07	57,57,57,57	0
3	CA	B	602	1/1	0.94	0.05	52,52,52,52	0
3	CA	A	601	1/1	0.97	0.04	32,32,32,32	0
3	CA	B	601	1/1	0.97	0.06	35,35,35,35	0
3	CA	C	601	1/1	0.98	0.06	32,32,32,32	0
3	CA	D	602	1/1	0.98	0.03	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GLC D 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.