



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 20, 2026 – 06:08 PM EDT

PDB ID : 9P19 / pdb\_00009p19  
Title : B. pseudomallei rubrerythrin room temperature structure  
Authors : Saha, S.; Budziszewski, G.R.; Russi, S.; Cohen, A.; Bowman, S.E.J.; Perry, S.  
Deposited on : 2025-06-09  
Resolution : 1.93 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>%</span> <span>92%</span> <span>7%</span> </div> </div>
1	B	140	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 97%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>%</span> <span>94%</span> <span>5%</span> </div> </div>
1	C	140	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 97%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>%</span> <span>97%</span> <span>..</span> </div> </div>
1	D	140	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 92%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>%</span> <span>92%</span> <span>7%</span> </div> </div>
1	E	140	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 98%);"></div> <div style="display: flex; justify-content: space-between; width: 90%; margin: 0 auto;"> <span>%</span> <span>98%</span> <span>..</span> </div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	140	<div><div>%</div><div><div></div></div><div>94%</div><div>6% •</div></div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12978 atoms, of which 6058 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rubrerythrin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	139	Total	C	H	N	O	S	0	2	0
			2106	674	1016	191	223	2			
1	B	139	Total	C	H	N	O	S	0	4	0
			2135	686	1028	194	225	2			
1	C	139	Total	C	H	N	O	S	0	0	0
			2072	668	994	189	219	2			
1	D	139	Total	C	H	N	O	S	0	1	0
			2096	672	1010	190	222	2			
1	E	139	Total	C	H	N	O	S	0	0	0
			2072	668	994	189	219	2			
1	F	139	Total	C	H	N	O	S	0	2	0
			2106	674	1016	191	223	2			

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Fe	0	0
			2	2		
2	B	2	Total	Fe	0	0
			2	2		
2	C	2	Total	Fe	0	0
			2	2		
2	D	2	Total	Fe	0	0
			2	2		
2	E	2	Total	Fe	0	0
			2	2		
2	F	2	Total	Fe	0	0
			2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	63	Total 64	O 64	0	1
3	B	75	Total 76	O 76	0	1
3	C	65	Total 65	O 65	0	0
3	D	52	Total 52	O 52	0	0
3	E	69	Total 70	O 70	0	1
3	F	52	Total 52	O 52	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Rubrerythrin



- Molecule 1: Rubrerythrin




- Molecule 1: Rubrerythrin




- Molecule 1: Rubrerythrin



- Molecule 1: Rubrerythrin



- Molecule 1: Rubrerythrin



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	203.36Å 203.36Å 69.87Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.35 – 1.93 29.35 – 1.93	Depositor EDS
% Data completeness (in resolution range)	95.3 (29.35-1.93) 94.5 (29.35-1.93)	Depositor EDS
$R_{merge}$	0.26	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.10 (at 1.93Å)	Xtriage
Refinement program	PHENIX 1.21.2_5419	Depositor
R, $R_{free}$	0.185 , 0.223 0.186 , 0.223	Depositor DCC
$R_{free}$ test set	3958 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.9	Xtriage
Anisotropy	0.573	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 42.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.013 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	12978	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/1112	0.64	0/1499
1	B	0.53	0/1129	0.69	0/1521
1	C	0.48	0/1100	0.62	0/1483
1	D	0.47	0/1108	0.58	0/1494
1	E	0.46	0/1100	0.57	0/1483
1	F	0.47	0/1112	0.57	0/1499
All	All	0.49	0/6661	0.61	0/8979

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	47	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1090	1016	1014	9	0
1	B	1107	1028	1036	6	0
1	C	1078	994	1006	2	0
1	D	1086	1010	1009	6	0
1	E	1078	994	1006	1	0
1	F	1090	1016	1014	4	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	A	64	0	0	0	0
3	B	76	0	0	0	0
3	C	65	0	0	0	0
3	D	52	0	0	0	0
3	E	70	0	0	0	0
3	F	52	0	0	0	0
All	All	6920	6058	6085	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 21 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:90:GLU:OE1	1:F:124:GLU:OE1	2.16	0.64
1:A:138:LEU:HD13	1:B:42[B]:ILE:CD1	2.30	0.61
1:D:90:GLU:OE1	1:D:124:GLU:OE1	2.24	0.56
1:A:103:LYS:HZ1	1:A:107:ASP:CG	2.14	0.55
1:A:138:LEU:HD13	1:B:42[B]:ILE:HD11	1.90	0.52

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/140 (99%)	137 (99%)	2 (1%)	0	100	100
1	B	141/140 (101%)	135 (96%)	6 (4%)	0	100	100
1	C	137/140 (98%)	134 (98%)	3 (2%)	0	100	100
1	D	138/140 (99%)	133 (96%)	5 (4%)	0	100	100
1	E	137/140 (98%)	135 (98%)	2 (2%)	0	100	100
1	F	139/140 (99%)	135 (97%)	4 (3%)	0	100	100
All	All	831/840 (99%)	809 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/106 (101%)	106 (99%)	1 (1%)	70	67
1	B	109/106 (103%)	108 (99%)	1 (1%)	70	67
1	C	105/106 (99%)	104 (99%)	1 (1%)	68	64
1	D	106/106 (100%)	105 (99%)	1 (1%)	70	67
1	E	105/106 (99%)	104 (99%)	1 (1%)	68	64
1	F	107/106 (101%)	106 (99%)	1 (1%)	70	67
All	All	639/636 (100%)	633 (99%)	6 (1%)	68	67

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	25	ARG
1	E	25	ARG
1	F	25	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	25	ARG
1	A	25	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	92	HIS
1	C	92	HIS
1	C	115	ASN
1	E	22	GLN
1	E	115	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	139/140 (99%)	-0.23	2 (1%) 73 79	12, 31, 66, 105	2 (1%)
1	B	139/140 (99%)	-0.22	1 (0%) 84 88	11, 31, 59, 84	4 (2%)
1	C	139/140 (99%)	-0.16	1 (0%) 84 88	23, 33, 61, 91	0
1	D	139/140 (99%)	-0.06	1 (0%) 84 88	21, 40, 69, 87	1 (0%)
1	E	139/140 (99%)	-0.15	1 (0%) 84 88	23, 34, 62, 79	0
1	F	139/140 (99%)	-0.01	2 (1%) 73 79	20, 41, 70, 94	2 (1%)
All	All	834/840 (99%)	-0.14	8 (0%) 79 84	11, 34, 68, 105	9 (1%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	2	ALA	4.6
1	E	2	ALA	4.0
1	F	2	ALA	3.9
1	A	67	VAL	3.8
1	B	2	ALA	3.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FE	D	202	1/1	0.69	0.15	34,34,34,34	1
2	FE	C	201	1/1	0.77	0.21	34,34,34,34	1
2	FE	A	201	1/1	0.79	0.13	32,32,32,32	1
2	FE	C	202	1/1	0.82	0.13	25,25,25,25	1
2	FE	F	201	1/1	0.83	0.10	31,31,31,31	1
2	FE	B	202	1/1	0.86	0.20	29,29,29,29	1
2	FE	E	201	1/1	0.87	0.11	39,39,39,39	1
2	FE	B	201	1/1	0.87	0.13	27,27,27,27	1
2	FE	A	202	1/1	0.88	0.09	38,38,38,38	1
2	FE	E	202	1/1	0.92	0.07	36,36,36,36	1
2	FE	D	201	1/1	0.92	0.06	29,29,29,29	1
2	FE	F	202	1/1	0.94	0.07	29,29,29,29	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.