



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 27, 2026 – 12:10 PM EDT

PDB ID : 9P3C / pdb\_00009p3c  
Title : Structure of radical S-adenosylmethionine methyltransferase, NocN, from *Nocardia* with SAH and side-ring closed product analog bound  
Authors : Wang, B.; Knox, H.L.; York, N.J.; Radle, M.I.; Silakov, A.; Booker, S.J.  
Deposited on : 2025-06-13  
Resolution : 1.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1365 (1.78-1.78)
Clashscore	190562	1395 (1.78-1.78)
Ramachandran outliers	187476	1382 (1.78-1.78)
Sidechain outliers	187428	1382 (1.78-1.78)
RSRZ outliers	180081	1365 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	
1	B	420	

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 7374 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called radical S-adenosylmethionine methyltransferase NocN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	386	Total	C	N	O	S	0	5	0
			3049	1930	538	567	14			
1	A	397	Total	C	N	O	S	0	6	0
			3135	1986	551	582	16			

- Molecule 2 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	Fe	S	0	0
			8	4	4		
2	A	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 3 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C<sub>14</sub>H<sub>20</sub>N<sub>6</sub>O<sub>5</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
3	A	1	Total	C	N	O	S	0	1
			52	28	12	10	2		

- Molecule 4 is 2-[(6S,13S)-6-[(2-{(1S,2R)-1-[(N-{2-[(1S)-1-amino-2-hydroxyethyl]-1,3-thiazole-4-carbonyl})-L-threonyl)amino]-2-hydroxypropyl}-1,3-thiazole-4-carbonyl)amino]-18-methyl-3,11,16-trioxo-1,3,4,5,6,11,12,13,14,16-decahydro-10,7-(azeno)-17,19-epimino-2,15,8,12-benzodioxathiazacycloicosin-13-yl]-1,3-thiazole-4-carboxylic acid (CCD ID: A1CG2) (formula: C<sub>42</sub>H<sub>44</sub>N<sub>10</sub>O<sub>13</sub>S<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total 69	C 42	N 10	O 13	S 4	0	0
4	B	1	Total 69	C 42	N 10	O 13	S 4	0	0
4	A	1	Total 69	C 42	N 10	O 13	S 4	0	0
4	A	1	Total 69	C 42	N 10	O 13	S 4	0	0

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is NICKEL (II) ION (CCD ID: NI) (formula: Ni).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ni	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	373	Total 373	O 373	0	0
7	A	392	Total 392	O 392	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: radical S-adenosylmethionine methyltransferase NocN



- Molecule 1: radical S-adenosylmethionine methyltransferase NocN



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.93Å 88.03Å 146.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.50 – 1.78 29.50 – 1.78	Depositor EDS
% Data completeness (in resolution range)	81.1 (29.50-1.78) 99.0 (29.50-1.78)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.27 (at 1.77Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487, PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.161 , 0.194 0.162 , 0.194	Depositor DCC
$R_{free}$ test set	2149 reflections (2.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.2	Xtriage
Anisotropy	0.110	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7374	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.77 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.7512e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH, CA, SF4, A1CG2, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.27	0/3218	0.47	0/4352
1	B	0.28	0/3127	0.48	0/4230
All	All	0.27	0/6345	0.47	0/8582

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3135	0	3119	15	0
1	B	3049	0	3023	14	0
2	A	8	0	0	0	0
2	B	8	0	0	0	0
3	A	78	0	57	0	0
3	B	52	0	38	0	0
4	A	138	0	0	3	0
4	B	138	0	0	4	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	392	0	0	0	0
7	B	373	0	0	0	0
All	All	7374	0	6237	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:408:LEU:HB2	4:B:504:A1CG2:N63	2.11	0.66
1:B:69:ASN:OD1	1:B:72:ARG:NH1	2.32	0.63
1:A:261:ARG:NE	1:A:271:GLU:OE1	2.33	0.58
1:A:399:ARG:HG3	1:A:400:LEU:HG	1.89	0.55
1:B:408:LEU:HB2	4:B:504:A1CG2:C56	2.37	0.54

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	401/420 (96%)	394 (98%)	7 (2%)	0	100	100
1	B	387/420 (92%)	381 (98%)	6 (2%)	0	100	100
All	All	788/840 (94%)	775 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/343 (96%)	330 (100%)	0	100	100
1	B	320/343 (93%)	320 (100%)	0	100	100
All	All	650/686 (95%)	650 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 3 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SAH	B	503	-	27,28,28	0.19	0	36,40,40	0.61	1 (2%)
2	SF4	A	501	1,3	0,12,12	-	-	-		
4	A1CG2	A	505	-	73,75,75	3.39	26 (35%)	104,108,108	2.89	46 (44%)
4	A1CG2	B	504	-	73,75,75	3.77	31 (42%)	104,108,108	2.92	47 (45%)
3	SAH	A	503[A]	2	27,28,28	0.15	0	36,40,40	0.34	0
4	A1CG2	A	504	-	73,75,75	3.24	18 (24%)	104,108,108	2.53	35 (33%)
3	SAH	A	502	-	27,28,28	0.28	0	36,40,40	0.59	1 (2%)
3	SAH	A	503[B]	2	27,28,28	0.17	0	36,40,40	0.48	0
3	SAH	B	502	2	27,28,28	0.17	0	36,40,40	0.63	1 (2%)
2	SF4	B	501	1,3	0,12,12	-	-	-		
4	A1CG2	B	505	-	73,75,75	3.93	34 (46%)	104,108,108	2.94	47 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SAH	B	503	-	-	1/15/31/31	0/3/3/3
2	SF4	A	501	1,3	-	-	0/6/5/5
4	A1CG2	A	505	-	-	19/79/81/81	0/6/7/7
4	A1CG2	B	504	-	-	6/79/81/81	0/6/7/7
3	SAH	A	503[A]	2	-	3/15/31/31	0/3/3/3
4	A1CG2	A	504	-	-	4/79/81/81	0/6/7/7
3	SAH	A	502	-	-	0/15/31/31	0/3/3/3
3	SAH	A	503[B]	2	-	2/15/31/31	0/3/3/3
3	SAH	B	502	2	-	2/15/31/31	0/3/3/3
4	A1CG2	B	505	-	-	18/79/81/81	0/6/7/7
2	SF4	B	501	1,3	-	-	0/6/5/5

The worst 5 of 109 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	505	A1CG2	C45-C48	23.50	1.71	1.48
4	A	504	A1CG2	C45-C48	22.85	1.70	1.48
4	B	504	A1CG2	C45-C48	22.50	1.70	1.48
4	A	505	A1CG2	C45-C48	22.43	1.70	1.48
4	B	504	A1CG2	C67-C62	6.76	1.50	1.39

The worst 5 of 178 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	505	A1CG2	C19-C21-N22	10.50	137.57	123.81
4	B	504	A1CG2	C19-C21-N22	9.17	135.83	123.81
4	A	504	A1CG2	C02-C03-N05	8.75	137.86	123.61
4	B	504	A1CG2	C02-C03-N05	8.62	137.65	123.61
4	A	505	A1CG2	C02-C03-N05	7.95	136.56	123.61

There are no chirality outliers.

5 of 55 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	502	SAH	C-CA-CB-CG
3	A	503[A]	SAH	O4'-C4'-C5'-SD
3	A	503[A]	SAH	C3'-C4'-C5'-SD
3	A	503[B]	SAH	O4'-C4'-C5'-SD
3	A	503[B]	SAH	C3'-C4'-C5'-SD


There are no ring outliers.

4 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	505	A1CG2	2	0
4	B	504	A1CG2	2	0
4	A	504	A1CG2	1	0
4	B	505	A1CG2	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


## Ligand SAH B 503




## Ligand SF4 A 501




## Ligand A1CG2 A 505



## Ligand A1CG2 B 504




## Ligand SAH A 503 (A)







## Ligand A1CG2 A 504




## Ligand SAH A 502





## Ligand SAH A 503 (B)



## Ligand SAH B 502







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	397/420 (94%)	-0.13	12 (3%)	52	60	9, 22, 45, 71	6 (1%)
1	B	386/420 (91%)	-0.17	11 (2%)	55	62	9, 23, 45, 76	5 (1%)
All	All	783/840 (93%)	-0.15	23 (2%)	53	61	9, 22, 45, 76	11 (1%)

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	413	ALA	4.8
1	A	399	ARG	3.3
1	A	232	THR	3.2
1	B	232	THR	2.9
1	A	242	ALA	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	A1CG2	B	505	69/69	0.77	0.16	21,56,118,128	0
4	A1CG2	B	504	69/69	0.83	0.13	24,50,87,104	0
4	A1CG2	A	505	69/69	0.85	0.13	14,46,61,68	0
3	SAH	B	502	26/26	0.90	0.10	21,27,40,43	0
4	A1CG2	A	504	69/69	0.95	0.09	13,23,54,63	0
3	SAH	A	503[A]	26/26	0.96	0.06	10,13,16,18	26
3	SAH	A	503[B]	26/26	0.96	0.06	10,14,17,28	26
3	SAH	A	502	26/26	0.96	0.06	12,16,19,20	0
3	SAH	B	503	26/26	0.97	0.05	12,17,23,26	0
5	CA	A	506	1/1	0.97	0.09	40,40,40,40	0
5	CA	B	506	1/1	0.98	0.08	32,32,32,32	0
2	SF4	B	501	8/8	0.98	0.04	18,21,21,22	0
6	NI	A	507	1/1	0.98	0.03	13,13,13,13	1
2	SF4	A	501	8/8	0.99	0.03	12,13,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around A1CG2 B 504:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around A1CG2 A 505:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SAH B 502:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around A1CG2 A 504:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SAH A 503 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around SAH A 503 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around SAH A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SAH B 503:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around SF4 B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around SF4 A 501:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.