



wwPDB EM Validation Summary Report ⓘ

Sep 14, 2026 – 01:30 pm BST

PDB ID : 9TU8 / pdb_00009tu8
EMDB ID : EMD-56252
Title : CryoEM structure of DruE:ATPgammaS:DNA from Druantia type III - dimer 1
Authors : Grass, L.M.; Himpich, S.; Hilal, T.; Loll, B.; Wahl, M.C.
Deposited on : 2026-01-08
Resolution : 3.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev133
Mogul : 1.8.4, CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8855 (3.40 - 4.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Y	2108	
1	y	2108	
2	I	30	
2	i	30	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	30	 50% 40% 10%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 34015 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DEAD/DEAH box helicase.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	Y	2068	Total	C	N	O	S	0	0
			16419	10354	2932	3061	72		
1	y	2050	Total	C	N	O	S	0	0
			16275	10265	2912	3026	72		

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Y	-3	GLY	-	expression tag	UNP A0A4P8BDA5
Y	-2	ALA	-	expression tag	UNP A0A4P8BDA5
Y	-1	GLU	-	expression tag	UNP A0A4P8BDA5
Y	0	PHE	-	expression tag	UNP A0A4P8BDA5
y	-3	GLY	-	expression tag	UNP A0A4P8BDA5
y	-2	ALA	-	expression tag	UNP A0A4P8BDA5
y	-1	GLU	-	expression tag	UNP A0A4P8BDA5
y	0	PHE	-	expression tag	UNP A0A4P8BDA5

- Molecule 2 is a DNA chain called DNA (5'-D(P*AP*CP*CP*AP*AP*AP*T)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
2	i	7	Total	C	N	O	P	0	0
			142	68	28	39	7		
2	I	27	Total	C	N	O	P	0	0
			551	262	101	161	27		


- Molecule 3 is a DNA chain called DNA (27-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	27	Total	C	N	O	P	0	0
			556	263	109	157	27		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
4	Y	5	Total	Zn	0
			5	5	
4	y	5	Total	Zn	0
			5	5	

- Molecule 5 is PHOSPHOTHIOPHOSPHORIC ACID-ADENYLATE ESTER (CCD ID: AGS) (formula: C₁₀H₁₆N₅O₁₂P₃S).





Mol	Chain	Residues	Atoms						AltConf
5	Y	1	Total	C	N	O	P	S	0
			31	10	5	12	3	1	
5	y	1	Total	C	N	O	P	S	0
			31	10	5	12	3	1	

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DEAD/DEAH box helicase







C1526	I1417	H1325	L1247	A1064	C982	M883	F798	V706	L611	ASP	F427	Y326
L1527	K1425	G1326	H1248	L1065	P983	G884	A799	S707	M612	VAL	I428	R327
S1528	K1425	V1329	L1250	K1066	T984	R886	D800	I708	E813	ASP	R429	Q328
C1529	S1432	T1251	L1147	D1067	T985	R887	F801	L711	I614	VAL	I329	VAL
G1530	T1433	T1251	L1148	S1068	R986	F887	C802	L712	I615	VAL	P330	P330
Q1537	T1434	Q1254	G1150	V1069	R987	R888	L803	M715	F821	TRR	A436	E334
P1538	E1438	L1255	K1151	L1072	A988	V889	Y804	Q727	F822	GLU	C437	E334
T1541	E1439	L1256	G1156	E1073	L989	K890	K812	Q727	S622	ASP	K440	K440
E1543	S1440	L1257	M1157	H1075	P991	M891	N813	D990	P623	GLU	Q441	Q441
N1544	R1345	T1259	G1161	H1075	L993	L892	N813	D990	P623	ASP	C442	C442
K1546	R1346	Q1260	K1161	L1079	L994	P895	T816	M715	F821	ASP	K445	L345
M1547	K1349	M1262	W1165	T1080	L1000	K897	E818	I731	E830	GLU	A446	L345
P1548	A1350	M1263	M1166	E1081	P1001	S819	L817	I732	Q631	LEU	H447	L348
E1549	A1351	Q1264	P1174	Q1082	G1002	P909	L820	I732	Q631	ASP	N450	K349
R1550	T1351	Q1265	Q1185	Q1083	S1003	G821	G821	E733	F835	ASN	V454	S352
E1551	GLN	Q1265	Q1185	N1084	G1003	Q910	L822	M735	N636	ASN	P455	P353
N1552	ALA	Q1265	Q1185	Q1085	S1004	T911	V823	S740	G837	SER	F456	Q354
T1553	ASP	Q1265	Q1185	Q1086	E1005	R912	H826	S740	F643	F355	G457	D355
Y1554	PRO	Q1265	Q1185	N1087	E1005	D913	Y827	H743	T642	R638	V459	L356
L1555	ASN	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L640	H362	H362
Y1556	GLY	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	Y363	Y363
L1557	GLY	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	C467	R370
R1558	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	D468	Q371
C1559	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	A471	V381
G1560	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	V473	R382
S1567	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	R478	A390
C1571	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	C479	R391
H1572	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	S480	W394
G1573	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	C482	W394
H1574	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	L487	L397
V1575	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	K490	H398
H1576	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	E491	H399
K1577	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	E492	L402
D1578	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	M493	L403
L1579	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	G495	L406
Y1583	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	D497	D497
S1584	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	L408	L408
S1585	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	M409	M409
V1590	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	S412	S412
L1594	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	R413	R413
L1601	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	P505	P505
I1602	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	ALA	ALA
R1603	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	ALA	ALA
D1604	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	GLU	GLU
F1605	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	VAL	VAL
E1605	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	ASP	ASP
V1606	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	PHE	PHE
A1607	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	ALA	ALA
A1608	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	ALA	ALA
R1609	GLU	Q1265	Q1185	Q1087	E1005	D913	Y827	H743	T642	L641	ALA	ALA



- Molecule 2: DNA (5'-D(P*AP*CP*CP*AP*AP*AP*T)-3')



- Molecule 2: DNA (5'-D(P*AP*CP*CP*AP*AP*AP*T)-3')



- Molecule 3: DNA (27-MER)



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	20914	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	42	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	96000	Depositor
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.260	Depositor
Minimum map value	0.000	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.012	Depositor
Recommended contour level	0.04	Depositor
Map size ($\text{\AA}$)	319.488, 319.488, 319.488	wwPDB
Map dimensions	288, 288, 288	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.1093334, 1.1093334, 1.1093334	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AGS, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	Y	0.11	0/16787	0.27	0/22755
1	y	0.10	0/16641	0.30	1/22555 (0.0%)
2	I	0.20	0/617	0.39	0/949
2	i	0.10	0/159	0.25	0/242
3	G	0.16	0/625	0.30	0/962
All	All	0.11	0/34829	0.29	1/47463 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	y	1643	TYR	CB-CA-C	-5.36	110.38	116.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Y	16419	0	16203	409	0
1	y	16275	0	16083	584	0
2	I	551	0	304	16	0
2	i	142	0	79	2	0
3	G	556	0	302	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	Y	5	0	0	0	0
4	y	5	0	0	0	0
5	Y	31	0	12	4	0
5	y	31	0	12	4	0
All	All	34015	0	32995	1004	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 1004 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:y:1529:CYS:SG	1:y:1554:HIS:CD2	2.58	0.96
1:y:1119:MET:HE1	1:y:1132:ARG:HG2	1.63	0.81
1:y:2074:LEU:HD13	1:y:2078:GLN:HB3	1.65	0.79
1:Y:1753:TYR:HA	1:Y:1841:ARG:HG3	1.66	0.76
1:Y:1967:VAL:HB	1:Y:2021:TRP:HE1	1.51	0.76

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Y	2060/2108 (98%)	1956 (95%)	104 (5%)	0	100	100
1	y	2042/2108 (97%)	1956 (96%)	86 (4%)	0	100	100
All	All	4102/4216 (97%)	3912 (95%)	190 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Y	1761/1795 (98%)	1733 (98%)	28 (2%)	55	69
1	y	1746/1795 (97%)	1713 (98%)	33 (2%)	50	66
All	All	3507/3590 (98%)	3446 (98%)	61 (2%)	52	68

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	y	57	VAL
1	y	1788	VAL
1	y	305	LEU
1	y	1735	VAL
1	y	2000	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	Y	1727	GLN
1	y	1636	GLN
1	y	13	ASN
1	y	1273	GLN
1	Y	1990	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 10 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	AGS	Y	2206	-	29,33,33	0.47	1 (3%)	39,52,52	0.66	1 (2%)
5	AGS	y	2206	-	29,33,33	0.46	1 (3%)	39,52,52	0.75	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	AGS	Y	2206	-	-	4/21/38/38	0/3/3/3
5	AGS	y	2206	-	-	6/21/38/38	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	y	2206	AGS	PG-S1G	2.15	1.95	1.90
5	Y	2206	AGS	PG-S1G	2.14	1.95	1.90

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	y	2206	AGS	PA-O3A-PB	-3.97	119.20	132.83
5	Y	2206	AGS	PA-O3A-PB	-3.31	121.48	132.83

There are no chirality outliers.

5 of 10 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	y	2206	AGS	PB-O3A-PA-O5'
5	y	2206	AGS	C5'-O5'-PA-O3A
5	y	2206	AGS	O4'-C4'-C5'-O5'
5	y	2206	AGS	C3'-C4'-C5'-O5'
5	y	2206	AGS	C4'-C5'-O5'-PA

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	Y	2206	AGS	4	0
5	y	2206	AGS	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-56252. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 144




Y Index: 144




Z Index: 144


6.2.2 Raw map



X Index: 144



Y Index: 144




Z Index: 144


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 149




Y Index: 132




Z Index: 174


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.04. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 571 nm³; this corresponds to an approximate mass of 516 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.256 $\text{\AA}^{-1}$

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.256 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.90	-	-
Author-provided FSC curve	3.97	4.58	4.02
Unmasked-calculated*	7.32	9.82	7.66

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.32 differs from the reported value 3.9 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-56252 and PDB model 9TU8. Per-residue inclusion information can be found in section [3](#) on page [6](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.04 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.04).

9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 95% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.04) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.9490	0.2890
G	0.8580	0.2390
I	0.9440	0.2980
Y	0.9680	0.3400
i	0.9300	0.3080
y	0.9340	0.2390

